

Case Of Osteum Secundum type Atrial Septal Defect Presenting As Postpartum Pulmonary Hypertension and Heart Failure with reduced EF

Dr. Siddharth Thakkar¹, Dr Charmi Kotak², Dr. Sunil Kumar³, Dr. Harshita Nouni⁴

¹Resident, Department of Medicine, SBKS MI & RC, Pipariya, Vadodara.

Email ID: sidthakker98@gmail.com

²Assistant Professor, Department of Medicine, SBKS MI & RC, Pipariya, Vadodara.

Email ID: Charmikotak06@gmail.com

³Professor, Department of Medicine, SBKS MI & RC, Pipariya, Vadodara.

Email ID: Sunilmdmedicine.pmch@gmail.com

⁴Resident, Department of Medicine SBKS MI & RC, Pipariya, Vadodara.

Email ID: harshitannouni28@gmail.com

Cite this paper as: Dr. Siddharth Thakkar, Dr Charmi Kotak, Dr. Sunil Kumar, Dr. Harshita Nouni, (2025) Case Of Osteum Secundum type Atrial Septal Defect Presenting As Postpartum Pulmonary Hypertension and Heart Failure with reduced EF. *Journal of Neonatal Surgery*, 14 (15s), 2247-2249.

ABSTRACT

Atrial septal defect (ASD) is one of the most common congenital heart diseases in adults. In many instances, it is diagnosed during or shortly after pregnancy. A 30 -year-old female with no known past medical history presented to the OPD with atypical chest pain and shortness of breath. She reported that she had delivered her third baby 6 months prior at an outside hospital and had been experiencing symptoms since delivery. Symptoms were noted after delivery at the outside hospital and she was referred to higher hospital. In the emergency department the patient was hypoxic at 94% on room air. Electrocardiogram showed right ventricle (RV) enlargement, LV strain pattern, P-pulmonale, and . Her EKG also had notching of the QRS in the inferior leads. A 2D echocardiogram displayed a large osteum-secundum atrial septal defect measuring 3.4 cm in diameter with a bidirectional atrial shunt. She was also noted to have a severely dilated right ventricle, and moderate tricuspid valve regurgitation with pulmonary artery pressure of 40 mmHg. Her ejection fraction was 15% and thin rim of pericardial effusion is seen. Pregnancy causes specific physiologic changes that affect hemodynamics and can unmask an ASD. Firstly, hormonal changes early in pregnancy decrease blood pressure by 5-10 mmHg, leading to an increase in cardiac output. Secondly, plasma volume can increase by up to 50% through the second trimester, also increasing cardiac output. These physiologic changes can exacerbate RV overload, especially in women with RV dilation and enlargement from untreated ASDs, leading to the characteristic symptoms of heart failure described above. Our case was different because the patient did not meet the strict criteria for surgical closure. There was concern that deferring surgical correction of the ASD would result in a net right to left shunt and subsequent Eisenmenger syndrome, a feared complication of an untreated ASD. After a discussion with the patient, the decision was made to proceed with conservative management with the hope of relief in signs and symptoms of ASD. Once diagnosed, the decision to close ASD is complicated, dependent on multiple factors, and requires adequate risk/benefit analysis, patient communication, and a multi-disciplinary approach.

1. INTRODUCTION

Atrial septal defect (ASD) is one of the most common congenital heart diseases in adults. In many instances, it is diagnosed during or shortly after pregnancy. The hemodynamic changes which occur during pregnancy can exacerbate underlying cardiac conditions, and in turn unmask a pre-existing condition. We present a case of a young, post-partum female who presented with symptoms of heart failure due to a large ASD.

2. CASE PRESENTATION

A 30 -year-old female with no known past medical history presented to the OPD with atypical chest pain and shortness of breath. She reported that she had delivered her third baby 6 months prior at an outside hospital and had been experiencing symptoms since delivery. Symptoms were noted after delivery at the outside hospital and she was referred to higher hospital. In the emergency department the patient was hypoxic at 94% on room air. Electrocardiogram showed right ventricle (RV) enlargement, LV strain pattern, P-pulmonale, and . Her EKG also had notching of the QRS in the inferior leads. Troponin

was negative, but A 2D echocardiogram displayed a large ostium secundum atrial septal defect measuring 3.4 cm in diameter with a bidirectional atrial shunt. She was also noted to have a severely dilated right ventricle, and moderate tricuspid valve regurgitation with pulmonary artery pressure of 40 mmHg. Her ejection fraction was 15% and thin rim of pericardial effusion is seen. She was discharged with diuretics and anti-platelets. Cardiologist was consulted regarding this and their opinion was to get TEE and RHC after stabilization of heart failure.

3. DISCUSSION

ASD is one of the most common congenital diseases in adults. It is important to diagnose patients early, as risk of morbidity and mortality increase if left untreated. The primary test to diagnose an ASD is a transthoracic echocardiogram (TTE). An EKG for an ostium secundum ASD may show “crochetage sign” (notching of R in the QRS complex of inferior leads), as seen in this case. ASD results in shunting of blood between systemic and pulmonary circulation. Initially shunting occurs from the left to right side of the heart, due to higher chamber pressures on the left side of the heart, with the degree of shunting directly correlated to the size of the ASD. This shunting leads to excessive blood flow in the RV and PA, and in turn causes RV and RA dilation. Over time, the dilation may result in RV failure, pulmonary hypertension, arrhythmias, and paradoxical embolism. The risk of pulmonary vascular disease increases with age and female gender. Symptoms of ASD are non-specific at time of presentation, and include chest pain, fatigue, and exertional dyspnea.

Pregnancy causes specific physiologic changes that affect hemodynamics and can unmask an ASD. Firstly, hormonal changes early in pregnancy decrease blood pressure by 5-10 mmHg, leading to an increase in cardiac output. Secondly, plasma volume can increase by up to 50% through the second trimester, also increasing cardiac output. These physiologic changes can exacerbate RV overload, especially in women with RV dilation and enlargement from untreated ASDs, leading to the characteristic symptoms of heart failure described above.

The postpartum period is also associated with significant hemodynamic changes. Within 10 minutes following vaginal delivery, cardiac output and stroke volume can increase by 60 to 70 percent. These changes most likely result from increased cardiac preload from auto transfusion of placental blood to the intravascular space. In addition, the decompression of inferior vena cava post-delivery results in increased preload. These changes can result in unmasking of a previously asymptomatic ASD.

AHA guidelines recommend closure of secundum ASD causing impaired functional capacity, RA or RV enlargement, and net left to right shunt sufficiently large to cause physiological sequelae ($Q_p/Q_s > 1.5$). Closure is recommended as long as systolic pulmonary artery pressure is less than 50% of systolic systemic pressure and pulmonary vascular resistance is less than 33% of the systemic vascular resistance. In more advanced disease with pulmonary hypertension, the ASD serves as a pop off valve. In cases where the pulmonary to systemic pressure ratio is more than 50%, closure of the ASD could lead to right sided pressure overload and is not recommended.

Treating an ASD in pregnancy depends on when it is diagnosed. Most women newly diagnosed with ASD during pregnancy can have closure deferred until six-months after delivery. In cases where an ASD is known in advance, closure is recommended before pregnancy. Further, if the known ASD has resulted in severe pulmonary hypertension, pregnancy is contraindicated as it can result in high morbidity and mortality for the mother and fetus. In general, pregnancy is well tolerated in most women with uncomplicated ASD.

Our case was different because the patient did not meet the strict criteria for surgical closure. There was concern that deferring surgical correction of the ASD would result in a net right to left shunt and subsequent Eisenmenger syndrome, a feared complication of an untreated ASD. After a discussion with the patient, the decision was made to proceed with conservative management with the hope of relief in signs and symptoms of ASD.

4. CONCLUSION

Atrial septal defect is one of the most common congenital heart disease in adults and can go undiagnosed into adulthood. It is important to diagnose patients early to prevent development of pulmonary hypertension and right ventricular failure. The hemodynamic changes during pregnancy and post-partum can unmask underlying, undiagnosed cardiac conditions, as in this case. Once diagnosed, the decision to close ASD is complicated, dependent on multiple factors, and requires adequate risk/benefit analysis, patient communication, and a multi-disciplinary approach.



Fig. 1

EKG showing crocheta sign with R wave notching



Fig. 2

2d echo showing Ostium secundum ASD

REFERENCES

- [1] Warnes CA, Williams RG, Bashore TM, et al. ACC/AHA 2008 Guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (writing committee to develop guidelines on the management of adults with congenital heart disease). *Circulation* 2008;118:e714-833. 10.1161/CIRCULATIONAHA.108.190690
- [2] Craig RJ, Selzer A. Natural history and prognosis of atrial septal defect. *Circulation* 1968;37:805-15. 10.1161/01.CIR.37.5.805
- [3] Regitz-Zagrosek V, Blomstrom Lundqvist C, Borghi C, et al. ESC Guidelines on the management of cardiovascular diseases during pregnancy: the Task Force on the Management of Cardiovascular Diseases during Pregnancy of the European Society of Cardiology (ESC). *Eur Heart J* 2011;32:3147-97. 10.1093/eurheartj/ehr218
- [4] Ouzounian JG, Elkayam U. Physiologic changes during normal pregnancy and delivery. *Cardiol Clin.* 2012 Aug;30(3):317-29. doi: 10.1016/j.ccl.2012.05.004. Epub 2012 Jun 20. PMID: 22813360
- [5] Beitzke D, Wolf F, Edelhauser G, Lammer J, Loewe C. Right heart dilatation in adult congenital heart disease: imaging appearance on cardiac magnetic resonance. *Br J Radiol.* 2011 Feb;84(998):188-93. doi: 10.1259/bjr/13711325. Epub 2010 Dec 15. PMID: 21159804; PMCID:PMC3473844.
- [6] Soma-Pillay P, Nelson-Piercy C, Tolppanen H, Mebazaa A. Physiological changes in pregnancy. *Cardiovasc J Afr.* 2016 Mar-Apr;27(2):89-94. doi: 10.5830/CVJA-2016-021. PMID: 27213856; PMCID:PMC4928162
- [7] Stout KK, Daniels CJ, Aboulhosen JA, Bozkurt B, Broberg CS, Colman JM, Crumb SR, Dearani JA, Fuller S, Gurvitz M, Khairy P, Landzberg MJ, Saidi A, Valente AM, Van Hare GF. 2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation.* 2019 Apr 2;139 (14):e698-e800. doi: 10.1161/CIR.0000000000000603. Erratum in: *Circulation.* 2019 Apr 2;139(14):e833-e834. PMID: 30586767.
- [8] Bredy C, Mongeon FP, Leduc L, Dore A, Khairy P. Pregnancy in adults with repaired/unrepaired atrial septal defect. *J Thorac Dis.* 2018 Sep;10(Suppl 24):S2945- S2952. doi: 10.21037/jtd.2017.10.130. PMID: 30305955; PMCID: PMC6174140