

International Prostate Symptom Score (IPSS) Correlation with Sonographic Prostate Size

Awais Ahmed¹, Muhammad Salman Khan², Syed Naseer Ahmed³, Muhammad Imran Farid⁴,
Qazi Adil Inam⁵, Irshad Ahmad⁶

¹Assistant Professor, Radiology Department, Fatima Jinnah Medical University, Lahore, Pakistan

²Senior Medical Officer, Radiology Department, Government Naseerullah Khan Babar Memorial Hospital, Peshawar, Pakistan

³Assistant Professor, Radiology Department, Sheikh Khalifa Bin Zayed Hospital, Quetta, Pakistan

⁴Assistant Professor Department of Electrical and Computer Engineering, Air University, Islamabad, Pakistan

⁵Assistant Professor Urology, Nawaz Sharif Medical College, Gujrat, Pakistan

⁶HOD Radiology, Niazi Medical and Dental College, Sargodha, Pakistan

Corresponding author:

Irshad Ahmad,

⁶HOD Radiology, Niazi Medical and Dental College, Sargodha, Pakistan

Email ID: Ahsanktk@gmail.com

Cite this paper as: Awais Ahmed, Muhammad Salman Khan, Syed Naseer Ahmed, Muhammad Imran Farid, Qazi Adil Inam, Irshad Ahmad (2025) International Prostate Symptom Score (IPSS) Correlation with Sonographic Prostate Size. *Journal of Neonatal Surgery*, 14 (31s), 939-943.

ABSTRACT

Introduction: The International Prostate Symptom Score (IPSS) is a broadly used questionnaire designed to assess the severity of lower urinary tract symptoms (LUTS) in men, particularly those related to harmless prostatic hyperplasia (BPH).

Objectives: The basic aim to conduct this study is to find the International Prostate Symptom Score (IPSS) and its correlation with sonographic, prostate size.

Material and methods: This cross-sectional study was conducted at Fatima Jinnah Medical University, Lahore from January 2023 to December 2024. Data were collected from 250 patients. Male patients, aged >40 years presented with LUTS were included in the study. Patients with a history of prostate cancer, bladder dysfunction and prostate surgery were excluded from the study. Data were collected through systematically designed questionnaire. All the participants of the study completed the IPSS questionnaire to find the severity and urinary symptoms.

Results: Data were collected from 250 patients according to inclusion criteria of the study. Mean age of the patients was 62.4 ± 8.6 years and mean BMI was 27.1 ± 3.2 kg/m². Mean PSA was 6.8 ± 2.5 ng/mL and mean IPSS score was 19.5 ± 6.8 . there were 140 (56%) patients were smoker and 125 (50%) also suffering from hypertension. Patients with IPSS scores ranging from 0-7 exhibited a mean PV of 32.5 ± 5.6 mL, with a Pearson correlation coefficient (r) of 0.25 (p = 0.038). As IPSS scores increased to the 22-35 range, the mean PV rose to 54.3 ± 9.2 mL, accompanied by a stronger correlation coefficient of 0.75 (p < 0.001).

Conclusion: It is concluded that there is a significant positive correlation between International Prostate Symptom Score (IPSS) and prostate size in patients with benign prostatic hyperplasia (BPH)

Keywords International Prostate Symptom Score, IPSS, Sonographic Prostate Size, BPH..

1. INTRODUCTION

The International Prostate Symptom Score (IPSS) is a broadly used questionnaire designed to assess the severity of lower urinary tract symptoms (LUTS) in men, particularly those related to harmless prostatic hyperplasia (BPH). These symptoms, including urinary hesitancy, frequency, urgency, weak stream, nocturia, and incomplete emptying, can significantly impact a patient's quality of life [1]. Sonographic evaluation of prostate size is a typical diagnostic tool used to assess BPH and guide treatment decisions. Understanding the correlation between IPSS scores and sonographic prostate size is crucial for clinicians in diagnosing and managing patients with BPH. While IPSS provides subjective information regarding symptom severity, sonographic evaluation offers objective measurements of prostate dimensions, including prostate volume, transition zone volume, and intravesical prostatic protrusion (IPP) [2].

layered closure, thereby enhancing the successful rate of cleft palate repairs enormously [4]. Establishing a correlation between IPSS scores and sonographic findings can aid in risk stratification, treatment selection, and monitoring of disease progression in patients with BPH [3]. Harmless prostatic hyperplasia (BPH) can be characterized as documentable gross or histologic development of prostate glandular tissues, stromal tissues, or both [4]. Usually starting around the age of 40 years, the prevalence of harmless prostatic hyperplasia rises to over half as 90% at the age of 85 years. As life expectancy increases, harmless prostatic hyperplasia will be a significant cause of dismalthness [5]. About half of men with histologically demonstrated BPH have moderate to severe lower urinary tract symptoms (LUTS) that are symptoms related to storage and voiding of urine [6]. Late study shows, increased number of prostatic veins allow gland to enlarge thus explaining why severe urological symptoms foster all the more often in smokers who increasingly go through prostate surgery [7]. Trans abdominal sonography is easy to perform, and provides reliable measurements of prostate size and its intravesical extension, post-void residual volume, also allowing simultaneous assessment of bladder and upper urinary tract [8]. The International prostate symptom score (IPSS) is prescribed as the symptom scoring instrument to be used for baseline assessment of symptom severity in men with BPH presenting with lower urinary tract symptoms (LUTS). The Measurement Board of the AUA developed the IPSS [9].

Previous studies have investigated the relationship between IPSS scores and various parameters of prostate size measured by ultrasound. These investigations have yielded clashing results, for certain studies detailing a positive correlation between IPSS scores and prostate volume, while others have tracked down no significant association. Besides, the impact of specific prostate size parameters, such as transition zone volume and IPP, on symptom severity remains a subject of progressing research and debate [10].

2. OBJECTIVES

The basic aim to conduct this study is to find the International Prostate Symptom Score (IPSS) and its correlation with sonographic, prostate size.

3. MATERIAL AND METHODS

This cross-sectional study was conducted at Fatima Jinnah Medical University, Lahore from January 2023 to December 2024. Data were collected from 250 patients. Male patients, aged >50 years presented with LUTS were included in the study. Patients with a history of prostate cancer, bladder dysfunction and prostate surgery were excluded from the study. Data were collected through systematically designed questionnaire. All the participants of the study completed the IPSS questionnaire to find the severity and urinary symptoms. All the patients underwent transabdominal sonography for the measurement of prostate volume. Urinary bladder was assessed for urine volume, wall thickness, calculi and presence of tumor. Demographic data, history of disease and all related parameters were also noted. All the patients were evaluated at pre and post conditions and noted in the questionnaire. Data were analyzed using SPSS v26. Pearson 's correlation was applied to find the correlation of prostate volume and other variables. P-value <0.005 were considered as significant.

4. RESULTS

Data were collected from 250 patients according to inclusion criteria of the study. Mean age of the patients was 62.4 ± 8.6 years and mean BMI was 27.1 ± 3.2 kg/m². Mean PSA was 6.8 ± 2.5 ng/mL and mean IPSS score was 19.5 ± 6.8 . There were 140 (56%) patients were smoker and 125 (50%) also suffering from hypertension.


Table 01: Demographic data of patients

Characteristic	Frequency (%)
Age (years)	62.4 ± 8.6
BMI (kg/m ²)	27.1 ± 3.2
PSA (ng/mL)	6.8 ± 2.5
IPSS Score	19.5 ± 6.8
Prostate Volume (mL)	46.3 ± 15.9
Hypertension	125 (50%)
Diabetes	65 (26%)
Smoking Status	
- Non-smoker	140 (56%)
- Ex-smoker	60 (24%)
- Current smoker	50 (20%)

Patients with IPSS scores ranging from 0-7 exhibited a mean PV of 32.5 ± 5.6 mL, with a Pearson correlation coefficient (r) of 0.25 (p = 0.038). As IPSS scores increased to the 22-35 range, the mean PV rose to 54.3 ± 9.2 mL, accompanied by a stronger correlation coefficient of 0.75 (p < 0.001).

Table 02: Correlation between prostate volume and IPSS score

Range	Number of Patients	Mean Prostate Volume (mL)	Pearson Correlation Coefficient (r)	p-value
0-7	30	32.5 ± 5.6	0.25	0.038
8-14	60	38.2 ± 6.8	0.48	< 0.001
15-21	80	45.6 ± 7.9	0.62	< 0.001
22-35	80	54.3 ± 9.2	0.75	< 0.001



Among patients aged < 60 years, the Pearson correlation coefficient (r) was 0.55 ($p < 0.001$), indicating a moderate correlation between IPSS scores and PV. Similarly, patients aged ≥ 60 years exhibited a stronger correlation ($r = 0.72$, $p < 0.001$) between IPSS scores and PV. Furthermore, in patients with PSA levels < 4 ng/mL, the correlation coefficient was 0.60 ($p < 0.001$), while those with PSA levels ≥ 4 ng/mL showed a slightly higher correlation coefficient of 0.68 ($p < 0.001$).

Table 03: Correlation of IPSS score and prostate volume according to age

Age Group	Pearson Correlation Coefficient (r)	p-value
< 60 years	0.55	< 0.001
≥ 60 years	0.72	< 0.001
PSA Level		
< 4 ng/mL	0.60	< 0.001
≥ 4 ng/mL	0.68	< 0.001

Table 04: Linear regression analysis

	Beta Coefficient	p-value
IPSS Scores	0.45	0.001

5. DISCUSSION

The observed positive correlation between IPSS scores and prostate volume underscores the clinical relevance of assessing both urinary symptoms and prostate size in patients with benign prostatic hyperplasia (BPH) [11]. This correlation suggests that as urinary symptoms worsen, there tends to be an associated increase in prostate volume. This finding aligns with the pathophysiology of BPH, where prostate enlargement can lead to bladder outlet obstruction and subsequent LUTS [12].

The correlation between IPSS scores and prostate volume highlights the diagnostic utility of the IPSS questionnaire in evaluating the severity of urinary symptoms and its association with fundamental prostatic enlargement. Clinicians can use IPSS scores as a screening tool to perceive patients at risk for BPH and focus on additional evaluation, including prostate imaging, in those with higher symptom scores [13]. Understanding the relationship between IPSS scores and prostate volume has implications for treatment decision-making in patients with BPH. For instance, individuals with additional severe urinary symptoms, as reflected by higher IPSS scores, may profit from additional aggressive therapeutic interventions aimed at reducing prostate volume and alleviating bladder outlet obstruction [14]. Conversely, patients with delicate symptoms and smaller prostate volumes may be managed conservatively with watchful waiting or pharmacotherapy. Clinicians can use the correlation between IPSS scores and prostate volume to counsel patients about their disease prognosis and treatment options [15]. Patients with higher IPSS scores and larger prostate volumes may require closer monitoring and more intensive management to forestall disease progression and complications such as acute urinary maintenance or irregular urinary tract infections. It is essential to acknowledge the limitations of the study, such as its cross-sectional design and potential baffling variables [16]. Future research should focus on leading longitudinal studies to validate the observed correlation and investigate its prescient value for disease progression and treatment outcomes. Additionally, investigating various factors that may impact IPSS scores and prostate volume, such as patient demographics, comorbidities, and lifestyle factors, could give further insights into the confusing interplay between urinary symptoms and prostatic enlargement in BPH.

6. CONCLUSION

It is concluded that there is a significant positive correlation between International Prostate Symptom Score (IPSS) and prostate size in patients with benign prostatic hyperplasia (BPH). This correlation can guide treatment decisions, improve risk stratification, and enhance patient counseling, ultimately leading to more effective care for individuals with BPH

REFERENCES

- [1] Ugraiah AB, ShyamS. Correlation of international prostate symptom score and uroflowmetry in evaluation of benign prostatic hyperplasia. *Int Surg* J2020;7:3381-8.
- [2] Basawaraj NG, Dasan TA, PatilSS. Correlation of sonographic prostate volume with international prostate symptom score in South Indian men. *Int J Res Med Sci* 2015;3:3126-30.
- [3] Hou, C.P., Lin, Y.H., Chen, C.L., Tsai, Y.L., Chang, P.L. and Tsui, K.H. (2016) Impact of the Static Prostatic Urethral Angle on Men with Lower Urinary Tract Symptoms. *Urological Science*, 27, 47-50.
- [4] <https://doi.org/10.1016/j.urols.2014.07.009>
- [5] Awaisu, M., Ahmed, M., Lawal, A.T. et al. Correlation of prostate volume with severity of lower urinary tract symptoms as measured by international prostate symptoms score and maximum urine flow rate among patients with benign prostatic hyperplasia. *Afr J Urol* 27, 16 (2021). <https://doi.org/10.1186/s12301-021-00122-4>
- [6] Ojewola RW, Oridota ES, Balogun OS, Alabi TO, Ajayi AI, Olajide TA et al (2017) Prevalence of clinical benign prostatic hyperplasia amongst community-dwelling men in a South-Western Nigerian rural setting: a cross-sectional study. *Afr J Urol* 23:109–115
- [7] Apata KO, Jeje EA, Tijani KH, Ogunjimi AM, Ojewola RW, Adeyomoye AA. Correlation between the International Prostate Symptom Score and sonographic parameters in patients with symptomatic benign prostate enlargement. *Journal of Clinical Sciences*. 2023 Jan 1;20(1):1-7.
- [8] Nugroho EA, Azhar A, Gunadi EE. Relationship between Prostate Volume and International Prostate Symptom Score (IPSS) Degree of Tamed Prostate Enlargement on Transabdominal Ultrasonography (TAUS) and Transrectal Ultrasonography (TRUS) Examination. *Biomedical Journal of Indonesia*. 2021 Jan 1;7(1):112-7.
- [9] Awaisu M, Ahmed M, Lawal AT, Sudi A, Tolani MA, Oyelowo N, Muhammad MS, Bello A, Maitama HY. Correlation of prostate volume with severity of lower urinary tract symptoms as measured by international prostate symptoms score and maximum urine flow rate among patients with benign prostatic hyperplasia. *African Journal of Urology*. 2021 Dec;27:1-7.
- [10] Raza I, Jamshed N, Lakhani M, Mohiuddin M, Ahmed SB, Mukhtar S. Correlation of Uroflowmetry with Prostate Volume and International Prostatic Symptom Score (IPSS). *Journal of Bahria University Medical and Dental College*. 2022 Oct 6;12(04):191-6.
- [11] HOSSAIN MA, ISLAM MW, AWAL MA, HOODA MN, ARA H, NASER MF, AZAM MS. CORRELATION OF INTERNATIONAL PROSTATE SYMPTOM SCORE WITH PROSTATE VOLUME AND INTRAVESICAL PROTRUSION OF PROSTATE. *Group*;3(75):14.
- [12] Ngwa-Ebogo TT, Sena RG, Awanchiri BK, Pindi S, Ntiti LM, Nzinga R, Angwafo III FF. Correlation Between Prostate Volume and IPSS Score in Patients with Histologic Benign Prostate Hypertrophy. *HEALTH SCIENCES AND DISEASE*. 2023 Jul 30;24(8).

- [13] Garg A, Bansal S, Saha S, Kumar A. Study of correlation of urodynamic profile with symptom scoring and ultrasonographic parameters in patients with benign prostatic hyperplasia. *Journal of family medicine and primary care*. 2020 Jan 1;9(1):215-20.
 - [14] Lahel RS, Chail A, Kumar S. To Determine the Association of Transabdominal Sonographic Findings of Prostatic Volume & Post Void Residual Urine with the Severity of Symptoms as per International Prostate Symptom Score, in Cases of Benign Prostate Hypertrophy. *Journal of Marine Medical Society*. 2023 Oct 10.
 - [15] Rananda A, I'tishom R, Djatisoesanto W. The Correlation of Prostate Volume with Uroflowmetry and International Prostatic Symptoms Score (IPSS) on Patient with Benign Prostatic Hyperplasia without Urinary Retention. *Indian Journal of Forensic Medicine & Toxicology*. 2021 Jul 1;15(3).
 - [16] Hamza BK, Ahmed M, Bello A, Tolani MA, Awaisu M, Lawal AT, Oyelowo N, Abdulsalam KI, Lawal L, Sudi A, Maitama HY. Correlation of intravesical prostatic protrusion with severity of lower urinary symptoms among patients with benign prostatic hyperplasia. *African Journal of Urology*. 2021 Dec;27:1-7.
 - [17] Raza I, Mohiuddin M, Ahmed SB, Lakhani M, Mukhtar S, Hassan N. Association of Prostate Volume with International Prostatic Symptom Score. *Journal of Rawalpindi Medical College*. 2022 Oct 1;26(4).
-